

## *Culex pipiens* (Diptera: Culicidae): A Bridge Vector of West Nile Virus to Humans

GABRIEL L. HAMER,<sup>1</sup> URIEL D. KITRON,<sup>2</sup> JEFFREY D. BRAWN,<sup>3,4</sup> SCOTT R. LOSS,<sup>4</sup>  
MARILYN O. RUIZ,<sup>2</sup> TONY L. GOLDBERG,<sup>2</sup> AND EDWARD D. WALKER<sup>5</sup>

J. Med. Entomol. 45(1): 125–128 (2008)

**ABSTRACT** Host-feeding patterns of *Culex pipiens* L. collected in southwest suburban Chicago in 2005 were studied using polymerase chain reaction (PCR) and DNA sequencing techniques. *Culex* spp. mosquitoes, most identified to *Cx. pipiens* and the remainder to *Cx. restuans* by PCR, had fed on 18 avian species, most commonly American robin (*Turdus migratorius*), house sparrow (*Passer domesticus*), and mourning dove (*Zenaida macroura*). Additional blood meals were derived from four mammal species, primarily humans and raccoons (*Procyon lotor*). During a West Nile virus (WNV) epidemic in 2005, West Nile virus (WNV) RNA was detected in heads and thoraces of five *Cx. pipiens* ( $n = 335$ , 1.5%) using quantitative PCR. The hosts of these virus-infected, blood-fed mosquitoes included two American robins, one house sparrow, and one human. This is the first report of a WNV-infected *Cx. pipiens* mosquito collected during an epidemic of WNV that was found to have bitten a human. These results fulfill a criterion for incrimination of *Cx. pipiens* as a bridge vector.

**KEY WORDS** *Culex pipiens*, West Nile virus, blood-meal analysis, bridge vector

West Nile virus (WNV) is now endemic throughout temperate North America, with annual amplification events and regular epizootics and epidemics. Although individuals of >60 mosquito species have tested positive for WNV (Centers for Disease Control, West Nile virus home page), species in the genus *Culex*, especially *Culex pipiens* L. in the eastern United States north of 36° latitude, have been implicated as the primary enzootic vectors, i.e., those responsible for transmission among bird reservoir hosts (Marra et al. 2004, Turell et al. 2005). More than 23 mosquito species have been implicated as potential bridge vectors or epidemic vectors, i.e., those responsible for transmission to humans (Marra et al. 2004, Turell et al. 2005). Recent, indirect evidence based on blood-meal analysis and theory suggests that *Cx. pipiens* serves as both an enzootic and an epidemic (i.e., “bridge”) vector (Apperson et al. 2004, Kilpatrick et al. 2005). The evidence includes data documenting *Cx. pipiens* feeding on both birds and mammals (Apperson et al. 2004) and results of a analytical risk model incorporating data on virus infection and feeding rates on birds, mammals, and humans suggesting that *Cx. pipiens* and

*Cx. restuans* are responsible for 80% of human WNV infection in the northeastern United States (Kilpatrick et al. 2005). However, empirical data showing a virus-infected mosquito biting a human has heretofore been lacking. Our objective was to use blood-meal analysis, individual mosquito virus infection detection, and molecular species identification methods to identify enzootic and epidemic mosquito vectors of WNV in an endemic transmission area in suburban Chicago, IL.

### Materials and Methods

We sampled blood-fed mosquitoes in southwest suburban Chicago in 2005 using gravid traps and aspirators at 15 study sites consisting of 11 residential neighborhoods, 3 cemeteries, and 1 wildlife refuge. This region has had a high incidence of human cases of West Nile viral meningoencephalitis since 2002 and of St. Louis encephalitis historically in 1975 (Ruiz et al. 2004). Blood-fed mosquitoes were processed individually to identify the blood-meal source, to identify the mosquito species, and to detect WNV. To identify the blood-meal source, we first scored the Sella stage of blood-meal digestion using an ordinal rating system (Detinova 1962) and digital photography. The abdomen was removed for blood-meal analysis and polymerase chain reaction (PCR) identification of the mosquito species, whereas the thorax and head were retained for RNA extraction and quantitative reverse transcriptase (RT)-PCR for virus detection. We amplified the vertebrate mitochondrial cytochrome B gene in the blood meal using four separate PCRs with established primer pairs, purified the amplicon (if

<sup>1</sup> Corresponding author: Department of Fisheries and Wildlife, Michigan State University, 13 Natural Resources Building, East Lansing, MI 48824 (e-mail: ghamer@msu.edu).

<sup>2</sup> Department of Pathobiology, University of Illinois, College of Veterinary Medicine, 2001 South Lincoln Ave., Urbana, IL 61802.

<sup>3</sup> Department of Natural Resources and Environmental Sciences, University of Illinois, 1102 South Goodwin Ave., Urbana, IL 61801.

<sup>4</sup> Department of Animal Biology, University of Illinois, 505 South Goodwin Ave., Urbana, IL 61801.

<sup>5</sup> Department of Microbiology and Molecular Genetics, Michigan State University, East Lansing, MI 48824.

**Table 1. Blood-meal analysis of mosquitoes collected in south-west suburban Chicago in 2005**

Host	Mosquito spp.		
	<i>Culex pipiens</i>	<i>Culex restuans</i>	<i>Culex</i> spp. <sup>a</sup>
Avian-derived blood meals (total)	191	27	5
American robin	69	13	4
Blue jay	10	2	
House sparrow	34	5	
Gray catbird	1	1	
House finch	14		
Common grackle	1		
European starling	2	2	
House wren	2		
American kestrel	2		
Northern cardinal	19	1	
Black-capped chickadee	2		
Cedar waxwing	1		
Cooper's hawk	1		
Mourning dove	30	2	1
American goldfinch	1		
Brown thrasher	1		
Swainson's thrush	1		
Mallard		1	
Mammal-derived blood meals (total)	55	8	2
Raccoon	9	2	1
Human	44	6	1
Domestic dog	1		
Gray squirrel	1		
NA <sup>b</sup>	89	7	4

<sup>a</sup> No PCR reaction.

<sup>b</sup> *Culex* mosquitoes that did not produce a PCR amplicon using the *Culex* spp. primer sets (*Cx. pipiens*, *Cx. restuans*, *Cx. salinarius*).

present), directly sequenced it, and compared the sequences to those in GenBank (Apperson et al. 2002, Cupp et al. 2004, Molaei et al. 2006). Results of negative controls were acceptable, and all positive controls (blood from 17 species of birds, 9 species of mammals, and 2 species of amphibians) were accurately and consistently identified to species. The same extracted DNA used for blood-meal analysis also was used for PCR-based molecular identification of *Culex* species to verify all morphological identifications (Crabtree et al. 1995). A quantitative RT-PCR method was used to detect WNV RNA in the head and thorax using empirically derived crossover thresholds to determine positive samples (Lanciotti et al. 2000, Hamer et al. 2007).

## Results

Of 398 blood-fed mosquitoes, most (84%) were identified as *Culex* spp. morphologically and as *Cx. pipiens* by PCR. Blood meals of 246 individual *Cx. pipiens* ( $n = 335$ , 73.4%) were identified successfully to an avian or mammalian host (Table 1). The success of identifying a blood meal was negatively correlated with the Sella score of the abdomen ( $r = -0.90$ ,  $df = 4$ ,  $P = < 0.01$ ). Based on the identified blood meals, *Cx. pipiens* fed primarily on birds ( $n = 191$ , 77.6%), with 18 species identified. The most common avian blood sources were American robin (*Turdus migratorius*;  $n = 86$ ), house sparrow (*Passer domesticus*;  $n = 39$ ), and mourning dove (*Zenaida macroura*;  $n = 33$ ). Mammal

**Table 2. Blood-fed mosquitoes collected in suburban Chicago, IL, that tested positive for WNV**

Date in 2005	Species	Blood-meal identification
July 19	<i>Cx. pipiens</i>	House sparrow
August 2	<i>Culex</i> spp. <sup>a</sup>	NA <sup>b</sup>
August 18	<i>Cx. pipiens</i>	American robin
August 19	<i>Cx. pipiens</i>	American robin
September 6	<i>Cx. pipiens</i>	Human
September 7	<i>Cx. pipiens</i>	NA
September 13	<i>Culex</i> spp. <sup>a</sup>	NA

<sup>a</sup> Mosquito identified morphologically as *Culex* spp. but did not produce a PCR amplicon using *Culex* primer sets (*Cx. pipiens*, *Cx. restuans*, *Cx. salinarius*).

<sup>b</sup> No PCR reaction.

feeding accounted for the remaining 22.4% of the *Culex* spp. blood meals, with humans ( $n = 51$ ) and raccoons (*Procyon lotor*;  $n = 12$ ) as the most common blood sources. Blood-meal analysis results for *Cx. restuans* revealed similar rates of avian feeding ( $n = 27$  or 35, 77%) and similar avian and mammalian host species as *Cx. pipiens*.

WNV RNA was detected in 5 of 335 blood-fed *Cx. pipiens* mosquitoes (1.5%) using quantitative RT-PCR (Table 2); two additional blood-fed *Culex* spp. mosquitoes tested positive for WNV but did not yield an amplicon after the blood-meal analysis PCR. The blood-meal host was identified in four of the five WNV-positive *Cx. pipiens* as follows: American robin ( $n = 2$ ), house sparrow ( $n = 1$ ), and human ( $n = 1$ ).

## Discussion

*Culex pipiens* is commonly considered to be ornithophilic (Marra et al. 2004, Turell et al. 2005), although the percentage of bird feeding varies substantially by region (84–96% feeding on birds in New York, Apperson et al. 2002, 2004; 93% in Connecticut, Molaei et al. 2006; 71% in Tennessee, but only 35% in New Jersey, Apperson et al. 2004; 87% in Maryland and Washington, DC, Kilpatrick et al. 2006; and 78% in this study). One possible explanation for this variation is substantial substructuring of *Cx. pipiens* populations, with mammal and bird feeding forms and hybrids, identified in Europe and New York (Fonseca et al. 2004, Kent et al. 2007). The structure of the population in metropolitan Chicago is not known, but our results show that the mammal feeding rate was comparatively high.

Samples from residential areas such as backyards of homes yielded 75% of the total *Culex* spp. mosquitoes in our study. Other recent blood-meal analysis studies with *Cx. pipiens* were done within urban areas, but actual sample sites were parks, uninhabited military forts, sewage treatment plants, golf courses, wood lots, and public thoroughfares (Apperson et al. 2002, 2004, Molaei et al. 2006). Collecting blood-fed mosquitoes in immediate proximity to human habitation could explain our finding of a high frequency of human feeding by *Culex* mosquitoes. Host availability was not quantified in this study; however, results of blood-meal

analysis should not be misconstrued as merely reflecting host preferences.

We found that 7 of 398 individual blood-fed mosquitoes were infected with WNV for an infection rate of 18 individuals per 1,000 (1.8%). Other research efforts conducted at the same sites in the same year found 227 positive pools of 1,195 tested, for an infection rate of 11 per 1,000 mosquitoes (1.1%) (Hamer et al. 2007). The Chicago area experienced a WNV epizootic and epidemic in 2005, during a drought, with Illinois ranking second in the United States in the total number of human WNV cases (Centers for Disease Control, West Nile virus home page). The high *Culex* spp. infection rate that reached 59 per 1,000 in late July (Hamer et al. 2007) provided an opportunity to identify virus-positive blood-fed mosquitoes and simultaneously to determine the origin of their blood meal. During peak transmission in July 2005, the probability that a WNV-infected *Culex* spp. mosquito fed on a human was 0.01 (i.e., WNV mosquito infection rate of  $0.059 \times$  human feeding rate of 0.173).

Incrimination of an epidemic vector of WNV requires demonstration of direct association of an infected vector with humans, a corollary of Koch's postulates. Our study contributes to the substantial evidence that *Cx. pipiens* serves as the enzootic vector of WNV in large parts of North America by showing that virus-infected mosquitoes feed on American robins and house sparrows. Moreover, our study implicates *Cx. pipiens* as a bridge or epidemic vector of WNV, because a mosquito with a disseminated virus infection in the head plus thorax was found to have fed on a human, the first recorded observation of such an event. Although the identity, infection, and clinical status of the bitten human are unknown, this finding adds support to the hypothesis that *Cx. pipiens* serves as both an enzootic and bridge vector. Our experience has been that WNV infection is rather common in *Culex* spp. and *Culex pipiens* in particular, but rare in non-*Culex* mosquitoes. In fact, we have found that non-*Culex* mosquitoes are typically rare during epidemics of WNV at our study site, whether infected or not. Mosquito testing efforts in 2005–2006 in the same region resulted in only seven positive non-*Culex* pools of 859 tested (an infection rate of 1 per 1,000 mosquitoes), those being one positive pool each of *Aedes vexans* (Meigen), *Anopheles quadrimaculatus* (Say), and *Ochlerotatus triseriatus* (Say) and two positive pools each of *Coquillettidia perturbans* (Walker) and *Ochlerotatus trivittatus* (Coquillett) (Hamer et al. 2007). In comparison, of 2,016 *Culex* spp. pools tested, 425 were positive (estimated infection rate by maximum likelihood method of 12 per 1,000 mosquitoes, a figure comparable to the infection rate of 18 per 1,000 reported here for individually tested mosquitoes). Although we cannot completely exclude the possibility of other mosquito species playing a role as bridge vectors in the Chicago metropolitan area, the finding of a virus-positive *Cx. pipiens* with a human-derived blood meal combined with the relatively high rate of human feeding by *Cx. pipiens* suggests that control efforts focused on *Cx. pipiens* alone may largely reduce both epizootic amplification and transmission risk to humans.

## Acknowledgments

We thank L. Abernathy, B. Bullard, M. Goshorn, J. McClain, M. Neville, B. Pultorak, and T. Thompson for providing assistance in the field and laboratory and W. Reisen, M. Turell, and C. Jones for valuable improvements to the manuscript. This research was supported by the NSF/NIH program in the Ecology of Infectious Diseases (Grant 04-29124).

## References Cited

- Apperson, C. S., B. A. Harrison, T. R. Unnasch, H. K. Hassan, W. S. Irby, H. M. Savage, S. E. Aspen, D. W. Watson, L. M. Rueda, B. R. Engber, and R. S. Nasci. 2002. Host-feeding habits of *Culex* and other mosquitoes (Diptera: Culicidae) in the Borough of Queens in New York City, with characters and techniques for identification of *Culex* mosquitoes. *J. Med. Entomol.* 39: 777–785.
- Apperson, C. S., H. K. Hassan, B. A. Harrison, H. M. Savage, S. E. Aspen, A. Farajollahi, W. Crans, T. J. Daniels, R. C. Falco, M. Benedict, M. Anderson, L. McMillen, and T. R. Unnasch. 2004. Host feeding patterns of established and potential mosquito vectors of West Nile virus in the eastern United States. *Vector Borne Zoonotic Dis.* 4: 71–82.
- Centers for Disease Control and Prevention. 2007. West Nile virus home page (<http://www.cdc.gov/ncidod/dvbid/westnile/index.htm>).
- Crabtree, M. B., H. M. Savage, and B. R. Miller. 1995. Development of a species-diagnostic polymerase chain-reaction assay for the identification of *Culex* vectors of St. Louis encephalitis-virus based on interspecies sequence variation in ribosomal DNA spacers. *Am. J. Trop. Med. Hyg.* 53: 105–109.
- Cupp, E. W., D. H. Zhang, X. Yue, M. S. Cupp, C. Guyer, T. R. Sprenger, and T. R. Unnasch. 2004. Identification of reptilian and amphibian blood meals from mosquitoes in an eastern equine encephalomyelitis virus focus in central Alabama. *Am. J. Trop. Med. Hyg.* 71: 272–276.
- Detinova, T. S. 1962. Age-grouping methods in Diptera of medical importance: with special reference to some vectors of malaria, vol. 47. World Health Organization, Geneva, Switzerland.
- Fonseca, D. M., N. Keyghobadi, C. A. Malcolm, C. Mehmet, F. Schaffner, M. Mogi, R. C. Fleischer, and R. C. Wilkerson. 2004. Emerging vectors in the *Culex pipiens* complex. *Science* 303: 1535–1538.
- Hamer, G. L., E. D. Walker, J. D. Brawn, S. R. Loss, M. O. Ruiz, T. L. Goldberg, A. M. Schotthoefer, W. M. Brown, E. Wheeler, and U. D. Kitron. 2007. Rapid amplification of West Nile virus: the role of hatch year birds. *Vector Borne Zoonotic Dis.* (in press).
- Kent, R. J., L. C. Harrington, and D. E. Norris. 2007. Genetic differences between *Culex pipiens f. molestus* and *Culex pipiens pipiens* (Diptera: Culicidae) in New York. *J. Med. Entomol.* 44: 50–59.
- Kilpatrick, A. M., L. D. Kramer, S. R. Campbell, E. O. Alleyne, A. P. Dobson, and P. Daszak. 2005. West Nile virus risk assessment and the bridge vector paradigm. *Emerg. Infect. Dis.* 11: 425–429.
- Kilpatrick, A. M., P. Daszak, M. J. Jones, P. P. Marra, and L. D. Kramer. 2006. Host heterogeneity dominates West Nile virus transmission. *Proc. R. Soc. Lond. B.* 273: 2327–2333.
- Lanciotti, R. S., A. J. Kerst, R. S. Nasci, M. S. Godsey, C. J. Mitchell, H. M. Savage, N. Komar, N. A. Panella, B. C. Allen, K. E. Volpe, B. S. Davis, and J. T. Roehrig. 2000. Rapid detection of West Nile virus from human clinical specimens, field-collected mosquitoes, and avian samples

- by a TaqMan reverse transcriptase-PCR assay. *J. Clin. Microbiol.* 38: 4066–4071.
- Marra, P. P., S. Griffing, C. Caffrey, A. M. Kilpatrick, R. McLean, C. Brand, E. Saito, A. P. Dupuis, L. Kramer, and R. Novak. 2004.** West Nile virus and wildlife. *Bioscience* 54: 393–402.
- Molaei, G., T. A. Andreadis, P. M. Armstrong, J. F. Anderson, and C. R. Vossbrinck. 2006.** Host feeding patterns of *Culex* mosquitoes and West Nile virus transmission, northeastern United States. *Emerg. Infect. Dis.* 12: 468–474.
- Ruiz, M. O., C. Tedesco, T. J. McTighe, C. Austin, and U. D. Kitron. 2004.** Environmental and social determinants of human risk during a West Nile virus outbreak in the greater Chicago area, 2002. *Int. J. Health Geogr.* 3: 11.
- Turell, M. J., D. J. Dohm, M. R. Sardelis, M.L.O. Guinn, T. G. Andreadis, and J. A. Blow. 2005.** An update on the potential of North American mosquitoes (Diptera : Culicidae) to transmit West Nile virus. *J. Med. Entomol.* 42: 57–62.

*Received 4 September 2007; accepted 11 September 2007.*

---