

Coexisting generalist herbivores occupy unique nutritional feeding niches

Spencer T. Behmer^{*†} and Anthony Joern[‡]

^{*}Department of Entomology, Texas A&M University, College Station, TX 77843-2475; and [‡]Division of Biology, Kansas State University, Manhattan, KS 66506-4901

Communicated by Perry L. Adkisson, Texas A&M University, College Station, TX, December 19, 2007 (received for review May 15, 2007)

A mainstay of ecological theory and practice is that coexisting species use different resources, leading to the local development of biodiversity. However, a problem arises for understanding coexistence of multiple species if they share critical resources too generally. Here, we employ an experimental framework grounded in nutritional physiology to show that closely related, cooccurring and generalist-feeding herbivores (seven grasshopper species in the genus *Melanoplus*; Orthoptera: Acrididae) eat protein and carbohydrate in different absolute amounts and ratios even if they eat the same plant taxa. The existence of species-specific nutritional niches provides a cryptic mechanism that helps explain how generalist herbivores with broadly overlapping diets might coexist. We also show that performance by grasshoppers allowed to mix their diets and thus regulate their protein-carbohydrate intake matched optimal performance peaks generated from no-choice treatments. These results indicate the active nature of diet selection to achieve balanced diets and provide buffering capacity in the face of variable food quality. Our empirical findings and experimental approach can be extended to generate and test predictions concerning the intensity of biotic interactions between species, the relative abundance of species, yearly fluctuations in population size, and the nature of interactions with natural enemies in tritrophic niche space.

biodiversity | competition | resource partitioning | physiological ecology | geometric framework

Ecological niches remain central to explaining community structure (1–7), despite some reservations (8). In general, for within-trophic level interactions, species that differ in their use of resources are more likely to coexist (9–11), even in highly variable environments (9, 10, 12–14). This “resource-partitioning” framework does not obviously apply to coexisting species that use essentially the same resources, including generalist herbivores that eat a diverse and broadly overlapping array of plants (3, 15–18). Alternate models are required to explain these patterns (7). However, because herbivores show differences in the blend of nutrients that maximize growth and fitness (19–21), an opportunity for niche diversification within a resource-partitioning framework exists at the level of macronutrient use rather than discrete plant taxa.

Ecologists search for assembly rules to explain community structure (22), usually as the combined outcome of top-down and bottom-up interactions (6, 7, 23). Although both are important in terrestrial herbivore communities, two observations argue for a critical importance of understanding bottom-up factors. First, interactions among herbivores leading to coexistence can be driven directly by limited food resources (24). Second, plant nutritional quality often mitigates top-down pressures (23, 25), although the converse may also be true. Generally speaking, herbivores are categorized as being specialists or generalists. Specialization is the case for $\approx 75\%$ of all plant-feeding insects (26), and plant secondary metabolites (PSMs) function as a primary assembly rule for coexistence among specialist plant-feeders because PSMs restrict membership in a specific feeding guild (27, 28). For insect herbivores, specializing on plants that

are unpalatable to other species effectively reduces potential interspecific competition. In contrast, PSM assembly rules do not readily predict coexistence of generalist herbivores that are adapted to eat a broad range of plant taxa. Instead, we suggest that community structure for generalist herbivores may be governed in part by assembly rules defined by differences in species-specific nutritional needs. Under this rule, the nutrients within the plants become the currency of importance, and competition or other biotic interactions among coexisting generalist herbivores would be mediated because each species occupies its own unique “nutritional niche.” Here, we define a nutritional niche as the blend and ratio of nutrients that maximize fitness (20, 29).

The extent to which coexisting generalist herbivores might occupy unique nutritional niches can be explored by using a recently developed experimental framework called the “geometric framework” (19–21) (henceforth GF). The GF is a state-space modeling approach that explores how an animal solves the problem of balancing multiple nutritional needs in a multidimensional and variable environment. It treats an animal as living within a multidimensional nutrient space where there are as many axes as there are functionally relevant (fitness-affecting) nutrients. There are >30 required nutrients for most animals, but protein and carbohydrates are among the most important for herbivores because their concentrations in plants are highly variable (depending on plant type, age, and growing conditions) and often limiting (26). The blend of protein and carbohydrate that the herbivore needs to ingest to optimize growth is called an “intake target,” and there is strong evidence that herbivores have evolved a suite of behavioral and physiological mechanisms that enable them to approach this point (19–21). For a generalist herbivore, the intake target can be reached by regulating the amount of an individual food eaten, eating from a range of different foods, or more likely, through a combination of these two mechanisms. A herbivore can reach its intake target by switching back and forth between two foods (represented as trajectories, or “rails” running through protein-carbohydrate “nutritional space”) that alone are nutritionally suboptimal but together are complementary [supporting information (SI) Fig. 3a]. This switching behavior represents a typical situation for most generalist herbivores. If food selection in generalist herbivores is viewed as a problem of collecting and regulating the intake of multiple nutrients, then the GF provides an experimentally accessible framework for explicitly testing whether species-specific nutritional niches and adaptive nutritional phys-

Author contributions: S.T.B. and A.J. designed research; S.T.B. and A.J. performed research; S.T.B. analyzed data; and S.T.B. and A.J. wrote the paper.

The authors declare no conflict of interest.

[†]To whom correspondence should be addressed at: Department of Entomology, Heep Building, Room 509, Texas A&M University, College Station, TX 77843-2475. E-mail: s-behmer@tamu.edu.

This article contains supporting information online at www.pnas.org/cgi/content/full/0711870105/DC1.

© 2008 by The National Academy of Sciences of the USA

Table 1. Diet pairings for each species plus MANOVA results comparing protein–carbohydrate consumption on the two treatments

Species	Diet combinations	df	Exact <i>F</i>	Prob > <i>F</i>
<i>M. angustipennis</i>	(a) p7:c35 + p35:c7	2, 13	1.79	0.205
	(b) p7:c35 + p28:c14			
<i>M. bivittatus</i>	(a) p7:c35 + p35:c7	2, 14	3.71	0.051
	(b) p7:c35 + p28:c14			
<i>M. differentialis</i>	(a) p7:c35 + p35:c7	2, 16	1.98	0.171
	(b) p7:c35 + p28:c14			
<i>M. femurrubrum</i>	(a) p7:c35 + p35:c7	2, 5	1.54	0.301
	(b) p7:c35 + p28:c14			
<i>M. flavidus</i>	(a) p7:c35 + p35:c7	2, 12	0.49	0.626
	(b) p7:c35 + p28:c14			
<i>M. foedus</i>	(a) p14:c28 + p35:c7	2, 9	0.79	0.482
	(b) p7:c35 + p28:c14			
<i>M. sanguinipes</i>	(a) p7:c35 + p35:c7	2, 16	0.51	0.612
	(b) p7:c35 + p28:c14			

Each species was tested on two treatments, and each treatment contained a pair of food dishes with different diets. The diets differed from one another in their protein (p) and digestible carbohydrate (c) content (e.g., p7:c35 = 7% protein and 35% carbohydrate, expressed on a dry mass basis). If each species is actively regulating protein–carbohydrate intake, no significant difference in protein–carbohydrate consumption should be observed.

ology can facilitate coexistence in generalist insect herbivore communities (SI Fig. 3 b–d).

We apply the GF to seven co-occurring North American grasshoppers within the genus *Melanoplus* that feed, at the population level (30), on a similar range of host plants, and we test the degree to which these coexisting species occupy unique nutritional niches. We also assess the ability of individuals to regulate nutrient intake to achieve a suitably balanced diet, and we briefly describe the buffering capabilities of diet regulation.

Results

We determined protein–carbohydrate intake targets for each species over their final nymphal stadium by allowing individuals of each species to mix diets between two food dishes that contained nutritionally imbalanced but complementary foods. There were two treatments for each species (Table 1), which allowed us to verify that protein–carbohydrate intake on a treatment was not simply a function of grasshoppers eating equally between the two food dishes (i.e., no discrimination). If grasshoppers actively regulate protein–carbohydrate intake, the amounts of protein and carbohydrate consumed in the two treatments should be similar. Without exception, no significant difference in the amounts of protein and carbohydrate eaten was observed between the two treatments for all species examined (Table 1). Having established that each species was able to regulate its protein–carbohydrate intake, we pooled the protein and carbohydrate consumption values across the two mixed treatments for each species. We then tested the extent to which protein–carbohydrate intake points among the seven species of grasshopper overlapped. Fig. 1 shows a bicoordinate plot of the mean amounts of protein and carbohydrate (\pm SEM) eaten by each species over the final nymphal stadium, and a comparison of the protein–carbohydrate intake points among the seven species revealed significant differences [multivariate ANOVA (MANOVA): $F_{12,198} = 19.68$, $P < 0.0001$]. Paired comparisons using contrasts were made between species that were “nearest neighbors” in nutritional space (Table 2). All comparisons except one species pair (*M. angustipennis* and *M. sanguinipes*) exhibited significant differences in terms of their protein–carbohydrate intake.

We also compared how insects that were free to mix diets and self-regulate protein–carbohydrate intake performed relative to grasshoppers that were restricted to single diets with different

protein:carbohydrate ratios. Our results show, with one possible exception (*M. flavidus*), that growth rate and development time were near optimal in grasshoppers that were free to mix (Fig. 2).

Discussion

This work documents the existence of an explicit nutritional physiological mechanism underlying patterns of food resource partitioning that can facilitate long-term species coexistence of generalist herbivores (2, 9, 10). Specifically, we show that coexisting *Melanoplus* species partition two key macronutrients, protein and digestible carbohydrates, by differentially regulating their intake (Fig. 1). This partitioning, expressed either in absolute amounts of protein and carbohydrate eaten (their intake targets) or as a protein:carbohydrate ratio (occupying an area, or wedge, of protein–carbohydrate nutritional space),




Fig. 1. Protein–carbohydrate intake targets (means \pm SEM) for each of the seven coexisting species. Each number in the figure corresponds to a particular species, and species that are statistically different from one another have different colored error bars [only species 1 and 7 (*M. angustipennis* and *M. sanguinipes*) were statistically similar; for details, see Table 1]. The gray dashed lines represent the two most extreme foods, p7:c35 and p35:c7, and the area between them represents the available nutrient space. The different colored solid lines define the range of nutrient space occupied by a particular species (as determined by the SEM of the intake targets). Overlap in nutrient space is only obvious for species 1 and 7. For a description of how nutrient space was determined for each species, see the *Materials and Methods*.

Table 2. Comparison of protein–carbohydrate consumption points for species that were nearest neighbors in nutritional space

Species comparison	df	Exact <i>F</i>	Prob > <i>F</i>
<i>M. bivittatus</i> (2) vs. <i>M. differentialis</i> (3)	2, 98	5.82	0.004
<i>M. bivittatus</i> (2) vs. <i>M. foedus</i> (6)	2, 98	7.46	0.001
<i>M. flavidus</i> (5) vs. <i>M. foedus</i> (6)	2, 98	10.09	<0.001
<i>M. flavidus</i> (5) vs. <i>M. sanguinipes</i> (7)	2, 98	7.52	0.004
<i>M. femurrubrum</i> (4) vs. <i>M. sanguinipes</i> (7)	2, 98	3.53	0.033
<i>M. angustipennis</i> (1) vs. <i>M. sanguinipes</i> (7)	2, 98	0.88	0.417
<i>M. angustipennis</i> (1) vs. <i>M. femurrubrum</i> (4)	2, 98	3.96	0.022

Protein–carbohydrate consumption points for species that were nearest neighbors in nutritional space were compared by using specified contrast statements after a significant overall species effect (MANOVA: $F_{12,198} = 19.68$, $P < 0.0001$). The number in parentheses after each species refers to the number assigned to that species in Fig. 1. All nearest-neighbor comparisons were significantly different from one another, except for *M. angustipennis* (1) vs. *M. sanguinipes* (7).

places coexisting species in clearly defined species-specific nutritional niches and may be an effective physiological mechanism to modulate the effects of interspecific competition during critical periods. These regulated species-specific intake targets have functional significance because they correspond to measured performance optima of developmental time and growth rate (Fig. 2). Inspection of performance data also indicates that some buffering capacity exists because relatively large changes in the protein:carbohydrate ratio of the diets result in only small performance penalties. This finding suggests the existence of a mechanism for dealing with often-encountered suboptimal food in a highly variable nutritional environment, although it may come with a cost. It is worth noting that the flatness of our performance curves across the more centrally located diets (e.g., p14:c28, p21:c21, and p28:c14), and to some extent the p35:c7 diet, may reflect the fact that we limited our work to a single developmental stage across a limited number of diets. Recently, Simpson *et al.* (31) advocated the construction of fitness landscapes over nutrient space, an approach that can reveal a sharp fitness peak at the intake target when a combined performance variable is calculated, for example, growth rate multiplied by percentage survival. In our work, survival was 100% in all but a few diets, so combining growth rate with percentage survival did not provide new insights.

For generalist insect herbivores, the availability of macronutrients such as protein and carbohydrate is highly variable within and among host plants, seasons, and years (32). When limiting, the relative concentrations of macronutrients among alternate food plants in a variable environment can reinforce species-specific niche diversification and determine the outcome of species interactions. For example, current theory of niche differentiation leading to long-term coexistence does not require that resources always be limiting or that species always compete with one another (2, 9). It is expected, however, that coexisting species will diverge in relevant niche space and that resources are sometimes limiting and result in competition, which appears to happen in grasshoppers (33, 34) and other insect herbivores (7). Additional field experiments that alter foliar-N levels directly affect grasshopper performance and species interactions in accordance with ideas presented here (25, 32). Within this framework, our approach provides a starting point for developing and testing specific predictions concerning competition among coexisting species with broadly overlapping diets, the relative abundance of coexisting species, and year-to-year population fluctuations. For example, *M. angustipennis* and *M. sanguinipes* share similar protein–carbohydrate intake points and occupy similar positions in nutritional space (their protein:carbohydrate intake ratio is $\approx 1:1$); competition between these two species should be particularly strong. These two species should also exist in similar numbers because they remove similar

absolute amounts of protein and carbohydrate from the environment. Additionally, their population sizes relative to the entire community should be relatively stable from year to year because they occupy a central location in protein–carbohydrate nutritional space that makes them less susceptible to yearly permutations in their nutritional environment. In contrast, competition between *M. femurrubrum* and *M. flavidus* should be weak because their protein–carbohydrate intake points, and ratios, are very different. However, because the protein–carbohydrate intake targets of these two species are located near the boundaries of nutritional space, yearly shifts in the nutritional environment may cause their populations to fluctuate in a more extreme manner from year to year. Exact predictions of locations of such boundaries in the nutritional landscape based on artificial diets might be tricky, however, because of structural attributes for packaging nutrients in plants and herbivore capabilities in extracting them; some nutrients become less available than others to herbivores (35). However, Jonas (36) has shown for *M. bivittatus* that the slopes of the boundaries do not change between artificial diets and ground plant material of equal nutritional quality, just the absolute amount of diet eaten, indicating that functional relationships remain intact.

The extent to which divergence in nutritional requirements by generalist insect herbivores can provide a mechanism for explaining species coexistence and fluctuations in relative abundance is largely unexplored. The GF provides an experimental framework that can inform us about ecological patterns among coexisting insect herbivores to explain patterns of community structure at some appropriate scale. In particular, insights regarding potential competitive interactions determined by bottom-up factors can be deduced. If food is limiting (24), diversification in nutrient space may also help explain the coexistence of a seemingly high diversity of generalist herbivores that share host plants in their diet.

Macronutrient availability is highly variable in time and space, which may contribute to coexistence over the short term, such as through a storage effect (2, 9). Theory concludes that resource partitioning is ultimately required to maintain species coexistence over the long term (9, 10). In this sense, however, we are not claiming that other factors do not contribute or are not important to the coexistence of generalist herbivores (37). Rather, our results provide an important underlying mechanism facilitating explanations based on population fluctuations (2, 38) or top-down effects of natural enemies (6, 7, 37, 39, 40) and extend their predictions. For example, diet selection in herbivores is greatly affected by recent encounters with predators (23, 25, 39), parasitoids (41), and pathogens (42) and often decreases the amount of food eaten and influences which plant species are selected and/or can shift protein–carbohydrate intake targets. In fact, such interactions make it more important to select food

34. Chase JM (1996) Differential competitive interactions and the included niche: An experimental analysis with grasshoppers. *Oikos* 76:103–112.
35. Clissold FJ, Sanson GD, Read J (2006) The paradoxical effects of nutrient ratios and supply rates on an outbreaking insect herbivore, the Australian plague locust. *J Anim Ecol* 75:1000–1013.
36. Jonas JL (2007) Nutrient resources and stoichiometry affect the ecology of above- and belowground invertebrate consumers. PhD dissertation (Kansas State Univ, Manhattan, KS), p 159.
37. Chase J, et al. (2002) The interaction between predation and competition: A review and synthesis. *Ecol Lett* 5:302–315.
38. Vandermeer JH (2006) Oscillating populations and biodiversity maintenance. *BioScience* 56:967–975.
39. Lima SL (1998) Stress and decision making under the risk of predation: Recent developments from behavioral, reproductive, and ecological perspectives. *Adv Study Behav* 27:215–290.
40. Holt RD, Lawton J (1994) The ecological consequences of shared natural enemies. *Annu Rev Ecol Sys* 25:495–520.
41. Bernays EA, Singer MS (2005) Taste alteration and endoparasites. *Nature* 436:476.
42. Lee KP, Cory JS, Wilson K, Raubenheimer D, Simpson SJ (2006) *Proc R Soc Lond, Ser B: Biol Sci* 273:823–829.
43. Holt RD (1977) Predation, apparent competition, and the structure of prey communities. *Theor Popul Biol* 12:197–229.
44. Berdegue M, Trumble JT, Hare JD, Redak RA (1996) Is it enemy free space? The evidence for terrestrial insects and freshwater arthropods. *Ecol Entomol* 21:203–217.
45. Mira A, Bernays EA (2002) Trade-offs in host use by *Manduca sexta*: Plant characters vs. natural enemies. *Oikos* 97:387–397.
46. Agrawal AA, et al. (2007) Filling key gaps in population and community ecology. *Front Ecol Environ* 5:145–152.
47. Agrawal AA, Lau JA, Hamback PA (2006) Community heterogeneity and the evolution of interactions between plants and insect herbivores. *Q Rev Biol* 81:349–376.
48. Vandermeer JH, Pascual M (2006) Competitive coexistence through intermediate polyphagy. *Ecol Complex* 3:37–43.
49. Mulkern GB, et al. (1969) *Food Habits and Preferences of Grassland Grasshoppers of the North Central Great Plains* (Agricultural Experiment Station, North Dakota State University, Fargo, ND regional publication No. 196).
50. Joern A (1985) Grasshopper dietary (Orthoptera: Acrididae) from a Nebraska sandhills prairie. *Trans Nebr Acad Sci* 8:21–32.
51. Behmer ST, Raubenheimer D, Simpson SJ (2001) Frequency-dependent food selection in locusts: A geometric analysis of the role of nutrient balancing. *Anim Behav* 61:995–1005.
52. Scheiner SM (2001) MANOVA: Multiple response variables and multispecies interactions. *Design and Analysis of Ecological Experiments*, eds Scheiner SM, Gurevitch J (Chapman & Hall, New York), pp 99–115.